

Roles of Antiferromagnetic Fluctuation in High Field Phase Diagram of Superconductors with Strong Paramagnetic Depairing

Ryusuke Ikeda and Kazushi Aoyama

Department of Physics, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan

Abstract

The high field phase diagram and magnetic properties of CeCoIn₅ below $H_{c2}(0)$ are examined from the picture regarding the high field and low temperature (HFLT) phase as a possible Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) vortex lattice. Crucial roles of antiferromagnetic (AFM) fluctuations enhanced close to $H_{c2}(0)$ are stressed. The FFLO vortex lattice with a longitudinal modulation parallel to the field is stabilized compared with those with lateral modulations as a consequence of the presence of AFM fluctuations. Further, an unusual field-induced enhancement of the flux distribution is argued to be a consequence of interplay between the paramagnetic depairing and AFM fluctuation, both of which are enhanced with increasing field.

Keywords: Heavy fermion superconductors; f-electron systems; vortex; FFLO superconductivity

1. Introduction

At present, it is believed that the high field and low temperature (HFLT) phase of the heavy fermion superconductor CeCoIn₅ with a d -wave pairing symmetry is a Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) vortex lattice with a paramagnetically induced modulation *parallel* to the applied magnetic field [1, 2]. This identification is based on several key observations [3, 4, 5] and their consistency with theoretical results [2]. For instance, the instability of HFLT phase via Cd or Hg doping [5], which induces a localized [6] antiferromagnetic (AFM) order, seems to imply a destruction of long range order of the FFLO modulation and is well understood if the HFLT phase includes an *inhomogeneous* AFM order [2]. In this sense, the FFLO picture of the HFLT phase of CeCoIn₅ is *compatible* with the recent observation of AFM order there [7]. Nevertheless, a relation between the HFLT phase and the AFM *fluctuation*, which is believed to be enhanced near $H_{c2}(0)$ [8], has not been well understood yet. Here, we will report on some results of our study on the vortex state taking account of both the strong paramagnetic depairing *and* the AFM quantum fluctuation enhanced near $H_{c2}(0)$.

2. Phase diagram in $\mathbf{H} \perp c$

As mentioned earlier [9], stability of the FFLO state with a modulation parallel to the field against those

with lateral modulations is ensured by going *beyond* the weak-coupling BCS approach and including some quasiparticle damping. In CeCoIn₅ with strong AFM fluctuation *near* $H_{c2}(0)$, this fluctuation-induced damping seems to be a main origin of making the states with *lateral* modulations relatively unstable, although the AFM fluctuation tends to suppress even the FFLO state with the longitudinal modulation (see Fig.1). On the other hand, the AFM fluctuation does not necessarily suppress the first order H_{c2} -transition, which is another peculiar feature in CeCoIn₅ in high fields [1, 3], and seems to also have a partial role of inducing the first order H_{c2} transition. An example of the H - T phase diagram in $\mathbf{H} \perp c$ we obtain in terms of a microscopically-derived Ginzburg-Landau (GL) free energy functional including the AFM fluctuation is shown in Fig.1. Since an applied pressure in real experiments corresponds to a reduction of AFM fluctuations, the dependences of the two (mean-field) transition curves and the onset of the first order H_{c2} -transition on the AFM fluctuation strength are consistent with the features found in experiments [4].

3. Flux distribution in $\mathbf{H} \parallel c$

Neutron scattering data on CeCoIn₅ in $\mathbf{H} \parallel c$ show [10] the vortex lattice form factor enhanced with increasing the field, which is an opposite trend to the




Figure 1: The resulting $H_{c2}(T)$ (dashed curves) and the FFLO transition curves (solid ones) in $\mathbf{H} \perp c$ for a smaller (red) and larger (black) strength of the AFM fluctuation. Each arrow indicates the onset of the first order H_{c2} -transition. The upturn of $H_{c2}(T)$ below $t = 0.2$ is an artifact of the expansion on the wavenumber of the FFLO modulation and can be improved. Here, h is the reduced magnetic field normalized by the orbital-limiting field in 2D case, and $t = T/T_{c0}$.

conventional one in type II superconductors. Although such an anomalous behavior has been explained in intermediate fields as a result of strong Pauli-paramagnetic effect [11], this interpretation is insufficient in higher fields where the AFM fluctuation is not negligible. If the damping effect suppressing the paramagnetic depairing is the main consequence of the AFM fluctuation, it seems difficult that the above-mentioned anomaly is attributed to the AFM fluctuation. However, the AFM fluctuation also leads to an effective increase of the electron correlation and thus, of the paramagnetic depairing, implying that the field dependence of roles of the AFM fluctuation is not intuitively clear. For this reason, we have investigated the flux density distribution of the vortex lattice based on the microscopically derived GL free energy. The flux distribution follows from the Maxwell equation. Expanding up to the lowest order in the squared pair field, $|\Delta|^2$, and also in a local value of AFM fluctuation correlator χ (> 0), we obtain the superconducting contribution to the longitudinal flux density parallel to the applied field in the form

$$B^{(s)}(\mathbf{r}) = 4\pi C_G |\Delta|^2 [\tilde{B}(\mathbf{r}) + \chi \tilde{B}^{\text{AF}}(\mathbf{r})], \quad (1)$$

where C_G is a scale factor measuring the magnitude of the local flux density in the familiar Abrikosov lattice near T_c . Figure 2 shows spacial dependences of $\tilde{B}(\mathbf{r})$ and $\tilde{B}^{\text{AF}}(\mathbf{r})$ on the coordinate between neighboring two vortices for the two cases of a small (dotted lines) and large (solid lines) Maki parameter α_M , which measures the Pauli-paramagnetic effect. The feature in $\tilde{B}$ that




Figure 2: Dependences of $\tilde{B}(\mathbf{r})$ (black) and $\tilde{B}^{\text{AF}}(\mathbf{r})$ (red) at $t = 0.1$ and $h = 0.1$ on the coordinates ranging from a vortex center ($\tilde{r} = 0$) to the midpoint ($\tilde{r} = 1.4$) between the two vortices. Here, a triangular lattice was assumed in a d-wave pairing. The dotted and solid curves correspond to those for $\alpha_M = 0.01$ and 6.5 , respectively.

the magnetic flux is accumulated near the vortex center with increasing the paramagnetic depairing is qualitatively the same as that in Ref.[11]. A crucial result in this study is that, as the red curves show, the fluctuation contribution $\tilde{B}^{\text{AF}}$ is competitive with $\tilde{B}$ for small Maki parameters, while it assists and *enhances* the flux accumulation in the vortex core induced by the paramagnetic effect for large enough α_M . It appears that, among roles of the AFM fluctuation, the mass enhancement of quasiparticles will overcome an increase of the quasiparticle damping. This enhanced flux accumulation due to the AFM fluctuation should be a direct origin of the unresolved enhancement [10] of the structure factor in higher fields and seems to imply an apparent presence of an AFM quantum critical point near $H_{c2}(0)$.

4. Acknowledgement

The work is supported by the Japan Society for the Promotion of Science.

- [1] A. Bianchi et al., Phys. Rev. Lett. **91** (2003) 187004.
- [2] R. Ikeda, Phys. Rev. Lett. **102** (2009) 069703 and references therein.
- [3] T. Watanabe et al., Phys. Rev. B **70** (2004) 020506(R); X.Gratens et al., arXiv: cond-mat/0608722.
- [4] C.F.Miclea et al., Phys. Rev. Lett. **96** (2006) 117001.
- [5] Y. Tokiwa et al., Phys. Rev. Lett. **101** (2008) 037001.
- [6] L.D. Pham et al., Phys. Rev. Lett. **97** (2006) 056404.
- [7] M. Kenzelmann et al., Science **321** (2008) 1652.
- [8] F. Ronning et al., Phys. Rev. B **71** (2005) 104528.
- [9] R. Ikeda, Phys. Rev. B **76** (2007) 134504.
- [10] A. Bianchi et al., Science **319** (2008) 177.
- [11] M. Ichioka and K. Machida, Phys. Rev. B **76** (2007) 064502.